Stroke Endovascular Thrombectomy (EVT) Protocol

QHC-Belleville General Hospital Emergency Transfer Guide for EVT For Weekday Cases only 07:30 to 14:30 h (Exclude Stat Holidays and Weekends)

Step 1: Determine if patient is candidate for rt-PA. If yes, then administer rt-PA as per protocol. Determine if patient is potential EVT candidate from Telestroke Neurologist consultation or QHC MRP (if not using Telestroke) after review of Multiphase CT Angiography (Multiphase CTA).

_	_		_				_
Sten	2: Review	Clinical I	Inclusion	and Exclu	ısion	Criteria	for FVT
JLCD							

Exclusion Criteria Inclusion Criteria No to all Below: Yes to all Below: Intubated □ Age 18 years or greater Severe or Fatal co-morbid illness □ NIH Stroke Scale greater than 5 Intracranial bleed ☐ Pre stroke functioning independently Non Contrast CT with ASPECTS greater than 5 Intracranial or Extra Cranial dissection Fibromuscular Dysplasia CTA with Intracranial large proximal artery occlusion in Difficult femoral access anterior circulation Severe Contrast Allergy Multiphase CTA with moderate to good collaterals **Step 3: Timeframe Guidelines** □ Patient can be transferred to KGH to arrive by 08:00 to 15:30 h □ Time from Stroke Onset to Groin Puncture (EVT start) at KGH should be less than 6 hours Aim for the following Time Targets: ☐ Time from patient arrival in BGH ED to time patient leaves BGH is less than or equal to 45 minutes Time from BGH CT (first slice) to KGH arrival time is less than 60 minutes **Step 4: Preparation for EVT Transfer** □ Confirm from Telestroke Neurologist (or MRP if Telestroke not used) that clearance was received from KGH Neurologist On Call for Stroke to proceed with EVT transfer If patient is receiving rt-PA: □ Arrange RN escort ☐ Start Drip and Ship Protocol □ Obtain EVT transfer kit Contact Hastings-Quinte Paramedic Service Duty Office Superintendent: 613-771-9366 extension 400 then hit "7" to arrange Code 4 Land Ambulance Transfer "Stroke Protocol" via dispatch If not already done & waiting for EMS crew (Never Delay Transfer to Complete): □ Draw CBC, lytes, urea, creatinine, INR, PTT, glucose, troponin, βHCG if female patient less than 50 years Start 1 IV of 0.9% NaCl & insert 1 Saline Lock preferably with an 18 Gauge needle in the right antecubital fossa unless contraindicated (If EMS crew on site, check if crew can insert 2 IVs en route) ☐ Remove clothes & send with patient □ Insert foley catheter Request 1 family member accompany EMS if possible. If not, obtain family contact number for transport team to inform the Stroke Team at KGH of any family contact details. Instruct family member to stay by the phone and keep line free Transfer with Ambulance Cardiac Monitor □ Fax relevant patient information to KGH ED including QHC ED Face Sheet

Step 5:

 Call KGH Emergency Department (ED). Ask to speak to the Charge Nurse and inform them you have a patient that meets the "EVT Protocol", whether or not on Drip & Ship Protocol with Nurse Escort, and the time patient left ED

FAX (613) 548-2420